

Pseudogap Phenomena and Phase Diagram in the 2-Band Hubbard Model

Akito KOBAYASHI¹, Atsushi TSURUTA¹, Tamifusa MATSUURA¹ and Yoshihiro KURODA^{1,2},

¹*Department of Physics, Nagoya University, Nagoya 464-8602*

²*CREST, Japan Science and Technology Corporation (JST)*

(Received)

High- T_c superconducting materials (HTSC) have anomalous properties such as the pseudo-gap or spin-gap *etc.*, in Hall coefficient, $1/T_1T$ and the density of states *etc.* First including effects of strong on-site repulsion between d-electrons at Cu-sites, we obtain quasi-particles with superexchange interaction J_s , whose band width tends to zero, *i.e.*, the system goes to insulator, as δ tends to zero. The quasi-particles correspond to Zhang-Rice singlet states. J_s larger than the band width combined with the 2-dimensional character of the system induces strong antiferromagnetic (AF) and superconducting (SC) fluctuations. We treat effects of the AF ones in the FLEX approximation and those of the SC ones in the self-consistent t -matrix approximation to show that both fluctuations in the under-doped region start to increase at T_0 as T decreases from $T \gg T_c$, the AF ones dominate the SC ones at $T > T_{sg}$, while SC ones dominate at $T < T_{sg}$. This cross-over of the fluctuations causes the anomalous phenomena in the under-doped region. We also obtain the $T - \delta$ phase diagram of HTSC consistent to one observed in experiments.

KEYWORDS: pseudogap, superconducting gap, superconductivity, superconducting fluctuation, spin fluctuation, superexchange interaction, Hubbard model

Since the discovery of the high- T_c superconducting materials (HTSC), the anomalous metallic phase have attracted much attention, where various anomalous properties have been observed in considerably wide temperature range higher than the superconducting transition temperature T_c ^{1,2,3,4,5,6,7,8}. The antiferromagnetic fluctuations (AF fluctuations) are observed in various experiments, *e.g.* the NMR¹, the neutron scattering² and the Hall coefficients^{3,4}. On the other hand, the pseudo-gap is also observed in various experiments, *e.g.* the tunneling spectroscopy⁵ and the angle-resolved photo emission spectroscopy (ARPES)^{6,7,8}, which is of the same size and has the same $\mathbf{k}$ -dependence as the superconducting gap (SC gap). There are several different views regarding the origin of the anomalous metallic phase, especially concerning the pseudogap. For example, the pseudogap may be interpreted as a spin-excitation gap in the singlet RVB states obtained in the t - J model^{9,10}, or as precursor of the SC gap^{11,12,13,14,15,16}.

Based on the above experimental results, we have suggested a mechanism of the superconductivity and the anomalous metallic phase¹³, where the AF fluctuations and the superconducting fluctuations (SC fluctuations) play essential roles as described in the followings. These fluctuations are induced by significant features of the HTSC, *i.e.* a quasi-2 dimensional metal in the vicinity of the Mott insulator. In metals near Mott insulators, doped carriers create narrow quasi-particle band inside the charge transfer gap called as in-gap states, which have band width proportional to the hole doping rate δ and interact with each other through the superexchange interaction J_s ^{17,18,19}. It is essentially the strong correlation effects. Strong AF fluctuations are induced by J_s

in the in-gap state. The pairing interaction mediated by the AF fluctuations induces the high- T_c superconductivity with $d_{x^2-y^2}$ symmetry. At the same time, the pairing interaction promotes strong SC fluctuations because of the quasi-2D electronic system, which induces the pseudogap in the quasi-particle spectrum. The pseudogap suppresses the AF fluctuations. Thus, effects of both of the AF fluctuations and the SC fluctuations need to be treated in a consistent way.

In our previous studies¹³, we have shown that the SC fluctuations induce the pseudogap and plays key roles in the anomalous metallic phase by using the self-consistent t -matrix approximation based on the in-gap state and J_s , which are derived in the 2-band Hubbard model with the help of the slave boson technique together with the $1/N$ -expansion theory^{17,18,19}. We have shown that $1/T_1T$ of the NMR increases as T decreases at high temperatures, and reaches a maximum at T_{sg} followed by a sharp drop in the under-doped region, due to the evolution of the pseudogap. The evolution is consistent with those of the ARPES experiments.⁸ It has been also shown that T_c is suppressed by the SC fluctuations in the under-doped region.

In the present study, we demonstrate the above scenario explicitly by taking the AF fluctuations in the fluctuation-exchange (FLEX) approximation and the SC fluctuations in the self-consistent t -matrix approximation to treat both fluctuations on the equal footing, which are also based on the in-gap state and J_s in the 2-band Hubbard model. By using the Green's functions thus obtained, we calculate T_c , the antiferromagnetic transition temperature T_N , $1/T_1T$, the spin susceptibility and the density of states. As a result, we obtain the T - δ

phase diagram which is consistent with that observed in HTSC.

First, we describe our model and formulation. We take the simplest version of the $U = \infty$ 2-band Hubbard model for describing the electronic system in the CuO_2 plane:

$$H = \varepsilon_p \sum_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \varepsilon_d \sum_{i\sigma} d_{i\sigma}^\dagger d_{i\sigma} + N_L^{-\frac{1}{2}} \sum_{i\mathbf{k}\sigma} \{t_{i\mathbf{k}} c_{\mathbf{k}\sigma}^\dagger d_{i\sigma} b_i^\dagger + h.c.\}, \quad (1)$$

which is treated within the physical subspace where local constraints $\hat{Q}_i = \sum_{\sigma} d_{i\sigma}^\dagger d_{i\sigma} + b_i^\dagger b_i = 1$ hold^{20, 21}. In the above, $c_{\mathbf{k}\sigma}$, $d_{i\sigma}$ and b_i are annihilation operators for a p -hole with wave vector $\mathbf{k}$ and spin σ , a d -hole at i -th site with spin σ and a slave boson at i -th site, respectively, and $t_{i\mathbf{k}} = t_{\mathbf{k}} \exp(-i\mathbf{k} \cdot \mathbf{R}_i)$, with $t_{\mathbf{k}} = 2t[1 - \frac{1}{2}(\cos k_x a + \cos k_y a)]^{\frac{1}{2}}$, where a is the lattice constant (we set $a = 1$), and N_L is the total number of lattice sites.

The quasi-particle Green's functions of the leading order in the $1/N$ -expansion are given by^{17, 18}

$$G_0(\mathbf{k}, \omega) = \sum_{\gamma=\pm} A_\gamma(\mathbf{k})/(\omega - E_\gamma(\mathbf{k}) + i0^+), \quad (2)$$

with $E_\gamma(\mathbf{k}) = \frac{1}{2}[\varepsilon_p + \omega_0 + \gamma((\varepsilon_p - \omega_0)^2 + 4bt_{\mathbf{k}}^2)^{1/2}]$, $A_\gamma(\mathbf{k}) = \gamma(E_\gamma(\mathbf{k}) - \omega_0)/(E_+(\mathbf{k}) - E_-(\mathbf{k}))$, where N represents the spin-orbital degeneracy of d -hole and then we set $N = 2$ in the present case. The binding energy ω_0 and the residue b of the pole in the slave-boson Green's function are self-consistently determined together with the chemical potential μ . Here, it is noted that the solution $E_-(\mathbf{k})$ denotes the in-gap state emerging inside the charge transfer gap, $\Delta \equiv \varepsilon_p - \varepsilon_d$, upon doping carriers to a Mott insulator. The band width and the residue of the in-gap state roughly equal $2\omega_0$ and b , respectively, and are proportional to δ in the under-doped region. The effects of the interactions among the in-gap states are of higher-order terms in the $1/N$ -expansion.^{22, 23, 24} Recently, it has been shown that quasi-particle interactions via the superexchange interaction J_s play dominant roles in under-doped systems.¹⁹

Now, we derive the coupled equations of the AF fluctuations and the SC fluctuations. The AF fluctuations via J_s are given by

$$V(\mathbf{q}, \omega) = J_s(\mathbf{q})/(1 - J_s(\mathbf{q})\tilde{\chi}^s(\mathbf{q}, \omega)), \quad (3)$$

with $J_s(\mathbf{q}) = -J_s(\cos(q_x) + \cos(q_y))$ and the irreducible spin susceptibility

$$\begin{aligned} \tilde{\chi}^s(\mathbf{q}, \omega) = & N_L^{-1} \sum_{\mathbf{k}} \Omega_{\mathbf{k}+\mathbf{q}, \mathbf{k}} \pi^{-1} \int dx f(x) \\ & \times [\text{Im}G(\mathbf{k} + \mathbf{q}, x)G(\mathbf{k}, -\omega + x)^* \\ & + G(\mathbf{k} + \mathbf{q}, \omega + x)\text{Im}G(\mathbf{k}, x) \\ & - \text{Im}F(\mathbf{k} + \mathbf{q}, x)F(\mathbf{k}, -\omega + x)^* \\ & - F(\mathbf{k} + \mathbf{q}, \omega + x)\text{Im}F(\mathbf{k}, x)], \end{aligned} \quad (4)$$

where $\Omega_{\mathbf{k}, \mathbf{k}'} = b^2 t_{\mathbf{k}}^2 t_{\mathbf{k}'}^2 / ((E_-(\mathbf{k}) - \omega_0)^2 (E_-(\mathbf{k}') - \omega_0)^2)$.

It was shown that a component with the $d_{x^2-y^2}$ sym-

metry among various components of the spin-fluctuation-mediated interaction contributes dominantly to the pairing interaction.²⁴ We take only the component with the $d_{x^2-y^2}$ symmetry of the pairing interaction as

$$v_d = N_L^{-2} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} \psi_{\mathbf{k}} V(\mathbf{k} - \mathbf{k}', 0) \psi_{\mathbf{k}'}, \quad (5)$$

with $\psi_{\mathbf{k}} = \cos(k_x a) - \cos(k_y a)$.

The pairing susceptibility $\tilde{\chi}^p(\mathbf{q}, \omega)$, which is the irreducible part of the t -matrix, is given by²⁵

$$\begin{aligned} \tilde{\chi}^p(\mathbf{q}, \omega) = & \tilde{\chi}_n^p(\mathbf{q}, \omega) \\ & + v_d \tilde{\chi}_a^p(\mathbf{q}, \omega) \tilde{\chi}_a^{p\dagger}(\mathbf{q}, \omega) / (1 - v_d \tilde{\chi}_n^p(\mathbf{q}, \omega)), \end{aligned} \quad (6)$$

where the pairing susceptibility due to the normal Green's functions is given by

$$\begin{aligned} \tilde{\chi}_n^p(\mathbf{q}, \omega) = & -N_L^{-1} \sum_{\mathbf{k}} \psi_{\mathbf{k}}^2 \Lambda_{\mathbf{k}, \mathbf{q}-\mathbf{k}} \pi^{-1} \int dx \\ & \times [f(x) \text{Im}G(\mathbf{k}, x)G(\mathbf{q} - \mathbf{k}, \omega - x) \\ & - f(-x)G(\mathbf{k}, \omega - x)\text{Im}G(\mathbf{q} - \mathbf{k}, x)], \end{aligned} \quad (7)$$

while that due to the anomalous Green's functions by

$$\begin{aligned} \tilde{\chi}_a^{p(\dagger)}(\mathbf{q}, \omega) = & -N_L^{-1} \sum_{\mathbf{k}} \psi_{\mathbf{k}}^2 \Lambda_{\mathbf{k}, \mathbf{q}-\mathbf{k}} \pi^{-1} \int dx \\ & \times [f(x) \text{Im}F^{(\dagger)}(\mathbf{k}, x)F^{(\dagger)}(\mathbf{q} - \mathbf{k}, \omega - x) \\ & - f(-x)F^{(\dagger)}(\mathbf{k}, \omega - x)\text{Im}F^{(\dagger)}(\mathbf{q} - \mathbf{k}, x)] \end{aligned} \quad (8)$$

and $\Lambda_{\mathbf{k}, \mathbf{k}'} = b^2 t_{\mathbf{k}}^2 t_{\mathbf{k}'}^2 / ((\pi T)^2 + \omega_0^2)^2$. In the above both $\Omega_{\mathbf{k}, \mathbf{k}'}$ and $\Lambda_{\mathbf{k}, \mathbf{k}'}$ are the vertices connecting p - and d -holes.¹⁹

Self-energy corrections due to the SC fluctuations in the t -matrix approximation are given by

$$\begin{aligned} \Sigma_{\text{SCf}}(\mathbf{k}, \omega) = & \psi_{\mathbf{k}}^2 N_L^{-1} \sum_{\mathbf{q}} \Lambda_{\mathbf{k}, \mathbf{q}-\mathbf{k}} \pi^{-1} \int dx \\ & \times [f(x)T(\mathbf{q}, x + \omega)\text{Im}G(\mathbf{q} - \mathbf{k}, x) \\ & - g(x)\text{Im}T(\mathbf{q}, x)G(\mathbf{q} - \mathbf{k}, x - \omega)^*], \end{aligned} \quad (9)$$

with the t -matrix of the SC fluctuations

$$T(\mathbf{q}, \omega) = v_d^2 \tilde{\chi}^p(\mathbf{q}, \omega) / (1 - v_d \tilde{\chi}^p(\mathbf{q}, \omega)), \quad (10)$$

where $g(E) = (\exp\{E/k_B T\} - 1)^{-1}$.

Self-energy corrections due to the AF fluctuations in the fluctuation exchange (FLEX) approximation are given by

$$\begin{aligned} \Sigma_{\text{AFf}}(\mathbf{k}, \omega) = & N_L^{-1} \sum_{\mathbf{q}} \Lambda_{\mathbf{k}, \mathbf{k}-\mathbf{q}} \pi^{-1} \int dx \\ & \times [g(x)\text{Im}V(\mathbf{q}, x)G(\mathbf{k} - \mathbf{q}, \omega - x) \\ & - f(-x)V(\mathbf{q}, \omega - x)\text{Im}G(\mathbf{k} - \mathbf{q}, x)]. \end{aligned} \quad (11)$$

The renormalized Green's function is given by

$$\begin{aligned} G(\mathbf{k}, \omega) = & [G_0(\mathbf{k}, \omega)^{-1} - \Sigma_{\text{SC}}(\mathbf{k}, \omega) \\ & - \Sigma_{\text{SCf}}(\mathbf{k}, \omega) - \Sigma_{\text{AFf}}(\mathbf{k}, \omega)]^{-1}, \end{aligned} \quad (12)$$

and the anomalous Green's function is

$$F(\mathbf{k}, \omega) = -G_n(\mathbf{k}, -\omega)^* \Delta_{SC}(\mathbf{k}) G(\mathbf{k}, \omega), \quad (13)$$

where the self-energy due to the SC gap

$$\Sigma_{SC}(\mathbf{k}, \omega) = -b^{-2} G_n(\mathbf{k}, -\omega)^* |\Delta_{SC}(\mathbf{k})|^2, \quad (14)$$

the renormalized Green's function without $\Sigma_{SC}(\mathbf{k}, \omega)$ by

$$G_n(\mathbf{k}, \omega) = [G_0(\mathbf{k}, \omega)^{-1} - \Sigma_{SCf}(\mathbf{k}, \omega) - \Sigma_{AFf}(\mathbf{k}, \omega)]^{-1} \quad (15)$$

and the SC gap with the $d_{x^2-y^2}$ symmetry by $\Delta_{SC}(\mathbf{k}) = v_d \psi_{\mathbf{k}} t_{\mathbf{k}}^2 \tilde{\Delta}_{SC}$. The order parameter $\tilde{\Delta}_{SC}$ is determined by

$$1 - \alpha = -v_d N_L^{-1} \sum_{\mathbf{k}} \psi_{\mathbf{k}}^2 \Lambda_{\mathbf{k}, \mathbf{k}} \pi^{-1} \int dx \times [f(x) \text{Im} G_n(\mathbf{k}, x) G(-\mathbf{k}, -x) - f(-x) G_n(\mathbf{k}, -x) \text{Im} G(-\mathbf{k}, x)]. \quad (16)$$

Both the SC and AF fluctuations are self-consistently treated by solving the coupled equations (3)-(16).

At $T = T_c$, eq. (16) corresponds to the Thouless criterion, $1 - v_d \tilde{\chi}^P(\mathbf{0}, 0) - \alpha = 0$, where α represents degree of the 3D effect in the quasi-2D electronic system. In the present model $1 - v_d \tilde{\chi}^P(\mathbf{0}, 0)$ does not reach zero at finite temperatures by the effect of $\Sigma_{SCf}(\mathbf{k}, \omega)$, because we take a 2D electronic system in the calculation of the above coupled equations. We take $\alpha = 0.5$ in the present study so as to have finite T_c of $O(100\text{K})$.

We define T_N as the temperature where $1 - J_s(\mathbf{Q}) \tilde{\chi}^s(\mathbf{Q}, 0) - \beta = 0$, where $\mathbf{Q} = (\pi, \pi)$ and β represents degree of the 3D effect. In the present model $1 - J_s(\mathbf{Q}) \tilde{\chi}^s(\mathbf{Q}, 0)$ also does not reach zero at finite temperatures by the effect of $\Sigma_{AFf}(\mathbf{k}, \omega)$ in the 2D electronic system. We take $\beta = 0.02$ in the present study.

By using the solutions of the above coupled equations, we calculate the NMR relaxation rate,

$$(T_1 T)^{-1} = N_L^{-1} \sum_{\mathbf{q}} F_{ab}(\mathbf{q}) \lim_{\omega \rightarrow 0} \text{Im} \chi^s(\mathbf{q}, \omega) / \omega, \quad (17)$$

where $\chi^s(\mathbf{q}, \omega) = \tilde{\chi}^s(\mathbf{q}, \omega) / (1 - J_s(\mathbf{q}) \tilde{\chi}^s(\mathbf{q}, \omega))$ and the factor $F_{ab}(\mathbf{q})$ is given in Ref. 26.

In actual numerical calculations, throughout the present study, we set $2t = 1.0$ (which is of $O(1\text{eV})$ in real systems), $\Delta = 2.5$ and $J_s = 0.1$. For an example, we have $\omega_0 = 0.0780$ and $b = 0.0515$ at $\delta = 0.1$. The total number of discrete points taken for the $\mathbf{q}$ -summation over the 2D first Brillouin zone is 32×32 . The ω -integral over the region from $-2\omega_0$ to $2\omega_0$ is replaced by the ω -summation of 80 discrete points.

Figure 1 shows the T - δ phase diagram obtained by solving the above coupled equations. Both the AF fluctuations and the SC fluctuations begin to increase at $T \cong T_0$ as T decreases. The AF fluctuations dominate in the region of $T_{sg} \lesssim T \lesssim T_0$, while the SC fluctuations dominate in the region of $T_c < T \lesssim T_{sg}$, where T_0 and T_{sg} increase monotonically when δ decreases. This crossover behavior induces the anomalous T -dependence of $1/T_1 T$ as discussed in Fig. 2. As a result of the domination by

the SC fluctuations in the under-doped region, T_c has a maximum at $\delta \cong 0.1$ and decreases as δ decreases. The key factor for obtaining these reasonable features is the narrow in-gap state with strong J_s , *i.e.* $J_s/W \gtrsim 1$ in the under-doped region, where the band-width W is approximately proportional to δ . This phase diagram accounts for essential features of the phase diagram observed in HTSC⁴⁾.




Fig. 1. The T - δ phase diagram obtained by solving the coupled equations (3)-(16). There is the superconducting state with $d_{x^2-y^2}$ symmetry in the region of $T \leq T_c$. The antiferromagnetic state exists in the region of $T \leq T_N$. The spin-gap temperature T_{sg} is defined as the temperature at which $1/T_1 T$ has a maximum as shown in Fig. 2. We define T_0 as the temperature at which the AF fluctuations begin to be appreciable, where explicit values of T_0 are determined by the condition that $[\frac{\partial}{\partial T}(1/T_1 T)|_{T=T_0}] / [(1/T_1 T)|_{T=0.08, \delta=0.1}] = -0.02$. We note that the SC fluctuations also become appreciable at $T \lesssim T_0$ as discussed in Fig. 3. The metallic region at $T \lesssim T_0$ is the anomalous metallic phase.

Figure 2 shows T -dependences of $1/T_1 T$ obtained by applying 3 different types of approximations for χ^s in eq. (17) at $\delta = 0.1$. In the case of RPA, $1/T_1 T$ diverges at $T = T_N^{\text{RPA}}$, which indicates the system becomes the antiferromagnetic state. In the case of FLEX, $1/T_1 T$ does not diverge but increases monotonically as T decreases. By solving the coupled equations including both the AF fluctuations and the SC fluctuations, $1/T_1 T$ is strongly suppressed in the low temperature region, and then has a maximum at T_{sg} . It is a typical example of the crossover behavior of the fluctuations in the anomalous phase.

Figure 3 shows ω -dependences of the normalized density of states $\rho(\omega)/\rho^0(\omega_0)$ at $\delta = 0.1$. The pseudogap begins to open at $T = 0.06 \cong T_0$ and develops gradually as T decreases, suppressing the AF fluctuations, which induces the crossover of the fluctuations. The pseudogap is induced by the SC fluctuations via the self-energy correc-




Fig. 2. T -dependences of $1/T_1 T$ obtained by applying 3 different types of approximations for χ^s in eq. (17) at $\delta = 0.1$. The solid line represents $1/T_1 T$ obtained by solving the coupled equations. The dashed line represents $1/T_1 T$ obtained by the FLEX approximation where only the self-energy corrections of the AF fluctuations are included. The dotted line represents $1/T_1 T$ obtained by the random phase approximation (RPA), *i.e.* no self-energy corrections of fluctuations are included.

tions, and has the same size and the same $\mathbf{k}$ -dependence as the SC gap, *i.e.* the precursor of the SC gap.




Fig. 3. ω -dependences of the normalized density of states $\rho(\omega)/\rho^0(\omega_0)$ at $\delta = 0.1$ and $T = 0.06, 0.05, 0.04, 0.03, 0.02$ and 0.01 , where $T_c = 0.012$. The density of states $\rho(\omega) = -\frac{1}{\pi N_L} \sum_{\mathbf{k}} \text{Im} G(\mathbf{k}, \omega)$ is obtained by solving the coupled equations, and $\rho^0(\omega)$ is obtained by using the Green's function of the in-gap state (eq. (2)) where no self-energy corrections of fluctuations are included.

The present study sheds light on a new problem in strongly correlated systems where two kinds of strong fluctuations affect each other. We have shown that the phase diagram of HTSC is described by our scenario consistently. The narrow in-gap state and strong J_s are the key factors for describing the features of the phase dia-

gram in the under-doped region. As for the quasi-particle interaction, the on-site Coulomb interaction is not effective in the under-doped region, because it is reduced to same order of the band-width of the in-gap state, while J_s remains to be finite almost independent of δ in the under-doping region¹⁹⁾.

The authors are grateful to Prof. M. Sato for fruitful discussions. The present work has been partially supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture, Japan.

-
- [1] H. Yasuoka, T. Imai and T. Shimizu: *Strong Correlation and Superconductivity*, eds. H. Fukuyama, S. Maekawa and A.P. Malozemoff (Springer-Verlag, Berlin, 1989) p. 254.
 - [2] L. P. Regnault, P. Bourges, P. Burlet, J. Y. Henry, R. J. Rossat-Mignod, Y. Sides and C. Vetter: *Physica B* **213-214** (1995) 48.
 - [3] T. Nishikawa, J. Takeda and M. Sato: *J. Phys. Soc. Jpn.* **63** (1994) 1441.
 - [4] M. Sato: *Physica C* **263** (1996) 271.
 - [5] Ch. Renner, B. Revaz, J. -Y. Genoud, K. Kadowaki and Ø. Fischer: *Phys. Rev. Lett.* **80** (1998) 149.
 - [6] A. G. Loeser, Z.-X. Shen, D. S. Dessau, D. S. Marshall, C. H. Parks, P. Fournier and A. Kapitulnik: *Science* **273** (1996) 325.
 - [7] H. Ding, T. Yokoya, J. C. Campuzano, T. Takahashi, M. Randeria, M. R. Norman, T. Mochiku, K. Kadowaki and J. Giapintzakis: *Nature* **382** (1996) 51.
 - [8] M. R. Norman, H. Ding, M. Randeria, J. C. Campuzano, T. Yokoya, T. Takeuchi, T. Takahashi, T. Mochiku, K. Kadowaki, P. Guptasarma and D.G. Hinks: *Nature* **392** (1998) 157.
 - [9] Y. Suzumura, Y. Hasegawa and H. Fukuyama: *J. Phys. Soc. Jpn.* **57** (1988) 2768.
 - [10] H. Fukuyama: *Prog. Theor. Phys. Suppl.* **108** (1992) 287.
 - [11] V. Emery and S. Kivelson: *Nature* **374** (1995) 434.
 - [12] Q. Chen, I. Kosztin, B. Jankó and K. Levin: *Phys. Rev. Lett.* **81** (1998) 4708.
 - [13] A. Kobayashi, A. Tsuruta, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **68** (1999) 2506; A. Kobayashi, A. Tsuruta, T. Matsuura and Y. Kuroda: *Physica B* **284-288** (2000) 661.
 - [14] S. Onoda and M. Imada: *J. Phys. Soc. Jpn.* **68** (1999) 2762.
 - [15] Y. Yanase and K. Yamada: *J. Phys. Soc. Jpn.* **68** (1999) 2999.
 - [16] S. Koikegami and K. Yamada: *J. Phys. Soc. Jpn.* **69** (2000) 768.
 - [17] H. Jichu, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **59** (1990) 2820.
 - [18] Y. Ōno, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **64**, (1995) 1595.
 - [19] S. Fukagawa, A. Kobayashi, K. Miura, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **67** (1998) 3536.
 - [20] P. Coleman: *Phys. Rev.* **B29** (1984) 3035.
 - [21] B. Jin and Y. Kuroda: *J. Phys. Soc. Jpn.* **57** (1988) 1687.
 - [22] K. Miura, T. Matsuura and Y. Kuroda: *Physica C* **179** (1991) 411.
 - [23] D.S. Hirashima, Y. Ōno, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **61** (1992) 649.
 - [24] M. Azami, T. Matsuura and Y. Kuroda: *J. Phys. Soc. Jpn.* **66** (1997) 2811.
 - [25] Y. Kosuge, H. Sato, A. Kobayashi, T. Matsuura and Y. Kuroda: *J. Phys. Chem. Sol.* in press.
 - [26] T. Imai: *J. Phys. Soc. Jpn.* **59** (1990) 2508.